

AD-A052 668

TEXAS A AND M UNIV COLLEGE STATION DEPT OF CHEMISTRY
NEW TRENDS IN THE CHEMISTRY OF ORGANOMETALLOPORPHYRINS. (U)

F/G 6/1

MAR 78 M TSUTSUI

N00014-75-C-0417

UNCLASSIFIED

TR-24

NL

1 OF 1
AD-A052 668



END
DATE FILMED
5-78
DDC



ADA052668

OFFICE OF NAVAL RESEARCH

Contract 15 N0014-75-C-0417
Task No. NR 053-559

R

(9) Technical Report No. 24

(6) New Trends in the Chemistry of Organometalloporphyrins.

by

(10) Minoru Tsutsui

Prepared for Publication

(14) TR-24

in the

Proceedings of VIIIth International Conference
On Organometallic Chemistry

Texas A&M University
Department of Chemistry
College Station, Texas 77843

(11) 15 Mar 78


(12) 10P.

March 15, 1978

AD No.
DDC FILE COPY

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited



407 997 708

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 24	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) New Trends in the Chemistry of Organometalloporphyrins	5. TYPE OF REPORT & PERIOD COVERED Interim	
7. AUTHOR(s) Minoru Tsutsui	6. PERFORMING ORG. REPORT NUMBER N00014-75-C-0417	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Texas A&M University College Station, Texas 77843	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-559	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217	12. REPORT DATE March 15, 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 8	
16. DISTRIBUTION STATEMENT (of this Report)	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
DISTRIBUTION STATEMENT A Approved for public release; Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted to Proceedings of VIIth International Conference on Organometallic Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) organometalloporphyrins metallocarbonylporphyrins out of plane metalloporphyrins dinuclear metalloporphyrins		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A broad review on current progress in chemistry of organometalloporphyrins is reviewed. An emphasis was placed on the development of metallocarbonylporphyrins, dinuclear metalloporphyrins and 'Skewered metalloporphyrins'.		


ABSTRACTION BY	
NTIS	WHM Series
SOS	DMT Series
<input checked="" type="checkbox"/> UNCLASSIFIED	
JOURNAL INFORMATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
DATA	AVAIL AND/OR SPECIAL
A	

NEW TRENDS IN THE CHEMISTRY OF ORGANOMETALLOPORPHYRINS

Minoru Tsutsui

Department of Chemistry, Texas A & M University, College Station, Texas 77843


Organometalloporphyrins in general are defined as synthetic metalloporphyrin complexes containing at least one direct metal-carbon bond. However, the coenzyme for vitamin B₁₂ (Figure 1) containing an adenosyl group linked to cobalt by a direct cobalt-carbon σ-bond is the first example of a naturally occurring organometallocorrin complex (similar to organometalloporphyrins in molecular structure) and the first-known stable organocobalt derivative.^{2,3} The determination of the structure of the vitamin B₁₂ coenzyme by x-ray crystallography^{4,5} prompted several studies of the synthesis and properties of cobalt-alkyl compounds.⁶⁻⁹ Johnson and his co-workers⁶⁻⁹ extended their studies to the porphyrin series by reaction of pyridinobromo-cobalt(III)actioporphyrin-I with a variety of alkyl- and aryl-magnesium halides in anhydrous 1,2-dimethoxyethane (DME) to give the corresponding alkyl- and aryl-cobalt(III) derivatives. Ethyl and p-tolyl-iron(III) derivatives¹⁰ of actioporphyrin-I were also prepared by a similar method. The first-row transition metal derivatives of organometalloporphyrins were thus synthesized for the first time. In an alternative synthesis, Johnson and his co-workers were also able to prepare the identical alkyl cobalt(III)actioporphyrin-I complexes as that mentioned above, by reduction of cobalt(II)actioporphyrin-I with 1% sodium amalgam in DME to form cobalt(I) species,¹¹ which then react with alkyl halides to give the final products. This method was also used for the preparation of hydroxylalkyl-cobalt(III)actioporphyrin-I complexes. Both the cobalt(III) and the iron(III) organometalloporphyrins are sensitive to light, especially in solution, and decompose by homolytic fission of the metal-carbon bond.



SPATIAL STRUCTURE OF COENZYME B₁₂

FIGURE 1

The use of metal carbonyls for the insertion of metal ions into porphyrins was first introduced by Tsutsui and his co-workers^{10,11} in 1966 (Figure 2). This method is probably one of the most important developments in porphyrin chemistry within the last two decades. In addition to a number of previously reported metallocoporphyrins, the reaction of metal carbonyls and metal carbonyl halides^{12,21,24-29} with neutral porphyrins has led to the synthesis of new metallocoporphyrin complexes of chromium, molybdenum, technetium, ruthenium, rhodium, rhenium and iridium.¹⁰⁻³² Except for the chromium and molybdenum complexes, carbonyl groups




USE OF $\text{Cr}(\text{CO})_6$ FOR INSERTION OF Cr^{II} TO PP

FIGURE 2

are retained by the metals in the new metallocporphyrin complexes.

By reaction of dirhenium decarbonyl, $\text{Re}_2(\text{CO})_{10}$, or ditechnetium decacarbonyl, $\text{Tc}(\text{CO})_{10}$, with mesoporphyrin IX dimethylester, $\text{H}_2\text{MPIXDME}$, in refluxing decalin under argon, Tsutsui^{1,3} and his co-workers have successfully prepared two unusual rhenium organometallocporphyrins ($\text{H}-\text{MP}[\text{Re}(\text{CO})]_2$), I, and $\text{MP}[\text{Re}(\text{CO})]_2$, II, a pair of technetium organometallocporphyrins,^{33,34} ($\text{H}-\text{MP}[\text{Tc}(\text{CO})]_2$), III, and $\text{MP}[\text{Tc}(\text{CO})]_2$, IV, and a mixed rhenium-technetium organometallocporphyrin, ($\text{OC}_2\text{ReMPtC}(\text{CO})_2$), V. A single crystal X-ray diffraction analysis of μ -[meso-tetr phenylporphinato]bis[tricarbonylrhenium(1)], TPP $[\text{Re}(\text{CO})]_2$, VI, (Figure 3), has shown



Porphyrin-Tridentate Ligand

Porphyrin-Hexadentate Ligand

FIGURE 3

that each rhenium ion is bonded to three nitrogen atoms and that two rhenium atoms are bonded to one porphyrin on opposite sides of the plane of the porphyrin molecule.

The metal ions in these complexes, I-VI, sit out of the plane of the porphyrin molecule. The monorhenium and monotechnetium complexes, I and III, where the porphyrin moiety acts as a tridentate ligand, resemble Fleischer's proposed "sitting-atop complex"^{35,39} and are good models for the intermediates in the insertion of a metal ion into porphyrin.⁴⁰ The dirhenium, ditechnetium, and mixed rhenium-technetium organometallocporphyrin complexes, II, IV, V, and VI, where the porphyrin moiety acts as a hexadentate ligand, are examples of the first isolated stable homo- and hetero-dinuclear organometallocporphyrin complexes.^{13,32} The monorhenium porphyrin complex, I, reacts with $\text{Re}_2(\text{CO})_{10}$ or $\text{Tc}_2(\text{CO})_{10}$ in refluxing decalin to form the dirhenium porphyrin complex,³¹ II, and the mixed rhenium-technetium porphyrin complex,¹³ V, respectively. Replacement of the pyrrolic proton ($\text{N}_{\text{H}}\text{H}^+$) of the monorhenium porphyrin complex by other metal ions such as Ag^+ , Mg^{2+} , and Pb^{2+} , has resulted in unstable complexes.³² The monotechnetium porphyrin complex, III, (Figure 4), behaves in a different manner by disproportionating to form a ditechnetium porphyrin complex, IV, and the free porphyrin, $\text{H}_2\text{MPIXDME}$, by heating in refluxing decalin. This unusual coordination phenomenon has never been reported. Such a reaction was not observed on heating monorhenium porphyrin complex, I, in refluxing decalin.^{31,32} It seems that both the rhenium and technetium dimetallocporphyrin complexes are thermodynamically more stable than the mono-metallocporphyrin complexes, because a reverse reaction of $\text{MP}[\text{M}(\text{CO})]_2$ to $(\text{H}-\text{MP})-\text{M}(\text{CO})_3$.

(M=Re or Tc), could not be detected between $M[\text{Ni}(\text{CO})_3]_2$ and $\text{H}_2\text{MPTCP}\text{H}_{3-3}$ in refluxing decalin for either the rhenium or technetium dimetalloporphyrin complexes.³⁴⁻³⁶





FIGURE 4

Temperature-dependent ^1H nmr spectral changes for $\text{HTPPRe}(\text{CO})_3$ and $\text{HDPRe}(\text{CO})_3$ dissolved in 1,1,2,2-tetrachlorethane showed fluxional behavior of $\text{Re}(\text{CO})_3$ group. This phenomenon is best explained by the intramolecular rearrangement of the metal carbonyl group among the four ring nitrogens of porphin and also movement of the N-H; it can also be regarded as an intramolecular substitution at rhenium or technetium, (Figure 4).³⁵

We have prepared a new salt type complex of porphyrins, monocation octaethylporphyrin tri- μ -halogeno-hexacarbonyldirhenate(I) from the reaction of $\text{Re}(\text{CO})_5\text{Cl}$ and H_2OEP in a 2:1 mole ratio in decalin. The structure of the complex was elucidated by the x-ray diffraction analysis method, (Figure 5).³⁶




STRUCTURE OF $(\text{H}_2\text{OEP})^+[\text{Re}_2(\text{CO})_6\text{Cl}_3]^- \text{H}_2\text{O}$

FIGURE 5

$[\text{Re}(\text{CO})_5\text{Cl}]_{2,\text{TPP}}$ has been oxidized by SbCl_5 in dichloromethane to yield $[\text{Re}(\text{CO})_3\text{Cl}]_{2,\text{TPP}} \cdot 2\text{SbCl}_5$ and $[\text{Re}(\text{CO})_3][\text{Re}(\text{CO})_3\text{Cl}] \cdot \text{SbCl}_5$. An x-ray determination of the structure of these complexes provides definite evidence for a "skewered complex", that is a metallocoporphyrin in which a metal-metal bond exists through the "hole" of the macrocycle, (Figure 6).³⁷

Two different methods were employed by Fleischer and his co-workers in preparing the rhodium and iridium porphyrin complexes.^{24,27} In one, the freshly prepared metal carbonyl halides, $[\text{Rh}(\text{CO})_3\text{Cl}]_2$ and $[\text{Ir}(\text{CO})_3\text{Cl}]_2$, were allowed to react with the porphyrins in glacial acetic acid solution to form the respective metallocoporphyrins, (Figure 7). In the second method,




First Skewered Complex (Radical Metalloporphyrin)

FIGURE 6

the cyclooctene complexes of rhodium and iridium were found to be reactive intermediates useful in the metalloporphyrin formation, (Figure 8). In both methods, incorporation of rhodium into the porphyrin was more readily achieved than was that of iridium. It is of interest that in the metalloporphyrins prepared by these methods, the iridium porphyrins retain the carbon monoxide ligand while rhodium and other metals do not.^{21,22} The carbon monoxide is very tightly bound to the iridium porphyrin moiety; heating, pumping, boiling in pyridine, or reprecipitating from concentrated sulfuric acid solution does not remove the carbonyl group from the complex.²⁴

By the reaction of $[\text{Rh}(\text{CO})_2\text{Cl}]$ ² with meso-tetraphenylporphyrin, H₂TPP, in refluxing benzene, two stable organometalloporphyrin derivatives of rhodium, Rh^{II}(CO)₂(TPP) · Cl and (c-phenyl)Rh^{II}-(TPP) · Cl, were separated by chromatography on an alumina column by Fleischer and his co-workers.^{21,25}

Yoshida and his co-workers were able to prepare two novel dinuclear rhodium(I) organometalloporphyrin complexes,^{26,27} VII and VIII, by modifying Fleischer's reaction conditions for the preparation of Rh^{II}(CO)(TPP) · Cl and (c-phenyl)Rh^{II}-(TPP) · Cl in refluxing benzene.^{22,23} Octaethylporphyrin, OEPH₂, or (N-methyl) octaethylporphyrin reacts with $[\text{Rh}(\text{CO})_2\text{Cl}]$, in benzene solution at room temperature under nitrogen atmosphere to produce VII and VIII. From the spectral data and the experimentally determined molecular weight,²⁹ VII was formulated as an acid, H⁺ [OEP · Rh^{II}(CO)₂Cl]⁻, which contains a Rh-Cl-Rh bridge. The proton NMR and infrared spectral data indicate that the $[\text{Rh}(\text{CO})_2\text{Cl}]$ moiety is maintained and the N-H and N-CH₃ bonds exist in VII. Since the Rh-Rh distance in $[\text{Rh}(\text{CO})_2\text{Cl}]$ ² has been reported to be 3.12 Å, and the distance between the two adjacent nitrogen atoms of planar porphyrin is about 2.9 Å, it was assumed²⁹ that the two Rh atoms of the $[\text{Rh}(\text{CO})_2\text{Cl}]$ moiety are bonded to the two adjacent nitrogen atoms of the porphyrinato core of VII, as



REACTION SCHEME FOR IRIDIUM PORPHYRIN
PREPARED VIA $\text{Ir}(\text{CO})_3\text{Cl}$

FIGURE 7




FIGURE 8





FIGURE 9

shown in Figure 9. Upon exposure to air, VII was slowly oxidized to form a rhodium(III) chloro complex of octaethylporphyrin, $\text{Rh}^{III}\text{Cl}(\text{OEP}) \cdot 2\text{H}_2\text{O}$, which can further react with alkyl lithium to give an alkyl-rhodium complex, another example of rhodium organometalloporphyrin complex. However, VIII behaves in a different manner to give the identical alkyl-rhodium complex either by gentle heating in chloroform or chromatography on silica gel (Figure 9). This phenomenon of alkyl migration from a nitrogen atom to a metal ion is reported for the first time. The alkyl migration may proceed concertedly with oxidation of rhodium(I) to rhodium(III). The $\text{N}-\text{CH}_3$ bond fission seems to be facilitated by the aid of a low-valent rhodium ion. The reaction of (N -ethyl)octaethylporphyrin with $[\text{Rh}(\text{CO})_2\text{Cl}]_2$ yields a rhodium(1) complex similar to VII , which is also easily oxidized to $\text{CH}_2\text{CH}_2 \cdot \text{Rh}^{III}(\text{OEP})$. The mechanism of metal oxidation and alkyl migration is still unknown.

Both rhenium carbonyl, $\text{Ru}_2(\text{CO})_{12}$, and ruthenium carbonyl halide, $[\text{Ru}(\text{CO})_2\text{Cl}_2]_2$, react with tetraphenylporphine to give the identical product, monocarbonyl-ruthenium(II) tetraphenyl porphine, TPPRuCO , IX . It was found that IX crystallizes with a molecule of either glycol or water and that these weakly bound molecules are bound to the carbonyl group. Recently, a single crystal x-ray diffraction analysis confirmed this structure. Imidazole and similar organic bases complex immediately at room temperature with TPPRuCO upon mixing in benzene. However, as expected for a low spin d system, substitution reactions of the monocarbonylruthenium(II) complex take place slowly and under severe conditions to replace the carbonyl group. Irradiation of degassed benzene or pyridine solutions of monocarbonyl ruthenium(II) antioporphyrin-I pyridinate with visible or ultraviolet light leads quantitatively to a ruthenium(II) porphyrin photodimer with a metal-metal bond. It is of interest that TPPRuCO in benzene solution reacts smoothly with excess nitric oxide to form a dinitrosyl-ruthenium(II) meso-porphyrin complex.


Since the successful isolation and purification of nitrogenase, the enzyme system that fixes molecular nitrogen in bacteria in living organisms, many studies have been made to find a nitrogenase model. Fleicher and co-workers created a model system consisting of the water soluble meso-tetra(p-sulfonatophenyl)porphinacobalt(III), $\text{Co}^{III}\text{TPPS}_4$, and sodium borohydride and found it to be very effective catalyst for the reduction of the substrate of nitrogenase. For example, acetylene was reduced catalytically in this system to ethylene and ethane. A possible mechanism for this catalytic system was proposed to include cobalt-acetylene and cobalt-ethylene metallocporphyrin π -complexes as intermediates (Figure 10). For the induced oxidation of cobalt(II) mesoporphyrin X dimethyl ester,



POSSIBLE MECHANISM FOR REDUCTION OF ACETYLENE BY A COBALT(III) PORPHYRIN SYSTEM.

FIGURE 10

$\text{Co}^{III}\text{TPPXYNE}$ by unsaturated hydrocarbons, an olefin-cobalt π -complex intermediate was proposed (Figure 11). Recently, the formation of an 1:1 adduct between cobalt(II) tetraphenylporphyrin, Co^{II}TPP , and carbon monoxide was reported by an esr study of a frozen



PROPOSED REACTION INTERMEDIATE FOR INDUCED OXIDATION
OF COBALT(II) PORPHYRIN BY CYCLOHEXENE

FIGURE 11

solution of toluene.⁴⁵ The paramagnetic low-spin d⁷ complex of Co^{II}TPP has a single unpaired electron in the d_{z²} orbital for σ bonding with carbon monoxide, which provides an opportunity for the examination by esr of σ spin delocalization to carbon monoxide. Similarly, the reversible binding of carbon monoxide to iron(II) protoporphyrin IX in piperidine was identified by its esr spectrum in frozen solution.

The σ-interaction in carbon monoxide to metalloporphyrin adducts and the proposed olefin-metal π-complex intermediates for induced oxidation of Cobalt(II) porphyrin by unsaturated hydrocarbon and the catalytic reduction of acetylene in Co^{II}TPPS and NaBH₄ model system has brought new examples of formation of organometalloporphyrins.

ACKNOWLEDGEMENT-This research was supported in part under a contract of the Office of Naval Research.

REFERENCES

1. Unusual Metalloporphyrins, XXX.
2. E.L. Smith, Vitamin B₁₂, 3rd Edition, Wiley, New York (1965).
3. G.N. Schrauser, Accounts Chem. Res., **1**, 97.
4. D.C. Hodgkin, Royal Society of London, Proceeding A, **288**, 294 (1965).
5. P.G. Lennert and D.C. Hodgkin, Nature, **192**, 937 (1961).
6. D. Dolphin and A.W. Johnson, Chem. Comm., 494 (1965).
7. D.A. Clarke, R. Grigg and A.W. Johnson, Chem. Comm., 208 (1966).
8. D.A. Clarke, R. Grigg, H. Pincock and A.W. Johnson, Chem. Comm., 309 (1967).
9. D.A. Clarke, D. Dolphin, R. Grigg, H. Pincock and A.W. Johnson, J. Chem. Soc. (C), **1**, 881 (1968).
10. M. Tsutsui, M. Ichikawa, F. Vohwinkle and K. Suzuki, J. Amer. Chem. Soc., **88**, 854 (1966).
11. M. Tsutsui, R. Velapoldi, K. Suzuki, F. Vohwinkle, M. Ichikawa and T. Koyano, J. Amer. Chem. Soc., **91**, 6262 (1969).
12. T.S. Srivastava and E.B. Fleischer, J. Amer. Chem. Soc., **92**, 5518 (1970).
13. M. Tsutsui and C.P. Hrung, J. Amer. Chem. Soc., **95**, 5777 (1973).
14. B.C. Chow and I.A. Cohen, Bioinorg. Chem., **1**, 57 (1971).
15. G.W. Sovocool, F. Hopf and D. Whitten, J. Amer. Chem. Soc., **94**, 4350 (1972).
16. J.J. Bonnet, S.S. Eaton, G.R. Eaton, R.H. Holm and J.A. Ibers, J. Amer. Chem. Soc., **95**, 2141 (1973).
17. T.S. Srivastava, L. Hoffman and M. Tsutsui, J. Amer. Chem. Soc., **94**, 1385 (1972).
18. M. Tsutsui, D. Ostfeld and L. Hoffman, J. Amer. Chem. Soc., **92**, 1820 (1971).
19. M. Tsutsui, D. Ostfeld, J.N. Francis and L. Hoffman, J. Coord. Chem., **1**, 115 (1971).
20. D. Cullen, E. Meyer, Jr., T. Srivastava and M. Tsutsui, J. Chem. Soc. Chem. Comm., 584 (1972).
21. E.B. Fleischer, R. Thorp and D. Venerable, Chem. Comm., 475 (1969).
22. J.W. Faller and J.W. Sibert, J. Organometal Chem., **31**, C5 (1971).
23. S.S. Eaton, G.R. Eaton and R. Holm, J. Organometal Chem., **39**, 179 (1972).
24. E.B. Fleischer and D. Lavallee, J. Amer. Chem. Soc., **89**, 7132 (1969).
25. N. Sadasivan and E.B. Fleischer, J. Inorg. Nucl. Chem., **30**, 591 (1968).
26. E.B. Fleischer and D. Lavallee, J. Amer. Chem. Soc., **89**, 7132 (1969).
27. B.R. James and D.V. Stynes, J. Amer. Chem. Soc., **94**, 6225 (1972).
28. Z. Yoshida, H. Ogoshi, T. Omura, E. Watanabe and T. Kuroasaki, Tetrahed. Lett., **11**, 1077 (1972).
29. H. Ogoshi, T. Omura and Z. Yoshida, J. Amer. Chem. Soc., **95**, 1666 (1973).
30. D. Cullen, E. Meyer, T.S. Srivastava and M. Tsutsui, J. Amer. Chem. Soc., **94**, 7603 (1972).
31. D. Ostfeld, M. Tsutsui, C.P. Hrung and D.C. Conway, J. Amer. Chem. Soc., **93**, 2548 (1971).
32. D. Ostfeld, M. Tsutsui, C.P. Hrung and D.C. Conway, J. Coord. Chem., **2**, 101 (1972).
33. M. Tsutsui and C.P. Hrung, Chem. Lett., **941** (1973).
34. M. Tsutsui and C.P. Hrung, J. Coord. Chem., **3**, 193 (1973).

35. N. Totsuji and C.P. Krung, *J. Amer. Chem. Soc.*, 96, 2630 (1974).
36. C.P. Krung, N. Totsuji, D. Cullen and E. Meyer, Jr., *J. Amer. Chem. Soc.*, 98, 7878 (1976).
37. S. Kato, N. Totsuji, *J. Amer. Chem. Soc.*, 99, 620 (1977).
38. E.B. Fleischer and J.H. Wong, *J. Amer. Chem. Soc.*, 82, 3493 (1960).
39. E.B. Fleischer, R.I. Choi, P. Hembright and A. Stone, *Inorg. Chem.*, 3, 1284 (1964).
40. R. Khodarpour and P. Hembright, *Inorg. Chem.*, 11, 133 (1972).
41. E.E. Van Tamelen, *Acc. Chem. Res.*, 3, 361 (1970).
42. G.H. Schrauzer and P.A. Domonay, *J. Amer. Chem. Soc.*, 93, 1606 (1971).
43. E.B. Fleischer and M. Krishnamurti, *J. Amer. Chem. Soc.*, 95, 1392 (1973).
44. N. Totsuji, A. Velapoldi, K. Suzuki and A. Ferrari, *J. Amer. Chem. Soc.*, 90, 2723 (1968).
45. B.R. Wayland and D. Mohajer, *J. Amer. Chem. Soc.*, 93, 5295 (1971).
46. D.V. Stynes, H.C. Stynes, B.R. James and J.A. Myers, *J. Amer. Chem. Soc.*, 95, 4087 (1973).

APPENDIX

TECHNICAL REPORT DISTRIBUTION LIST

<u>No. Copies</u>	<u>No. Copies</u>		
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market St., Rm. 447 San Francisco, Calif. 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Dr. R. M. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1
Commander Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1
		Dr. C. Quicksall Georgetown University Department of Chemistry 37th & O Streets Washington, D.C. 20007	1

TECHNICAL REPORT DISTRIBUTION LIST

<u>No. Copies</u>		<u>No. Copies</u>	
1	Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019
1	Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802
1	Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916		
1	Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712		
1	Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514		
1	Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139		
1	Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540		
1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154		
1	Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201		